arXiv:1411.6963v1 [math.NT] 14 Nov 2014

On certain quaternary quadratic forms

By Kazuhide Matsuda

Department of Engineering Science, Niihama National College of Technology,
7-1 Yagumo-chou, Niihama, Ehime, Japan, 792-8580
E-mail: matsuda@sci.niihama-nct.ac.jp
Fax: 0897-37-7809

Abstract In this paper, we determine all the positive integers a, b and c such that every nonnegative integer can be represented as

$$f_c^{a,b}(x,y,z,w) = ax^2 + by^2 + c(z^2 + zw + w^2) \text{ with } x, y, z, w \in \mathbb{Z}.$$

Furthermore, we prove that $f_c^{a,b}$ can represent all the nonnegative integers if it represents $n = 1, 2, 3, 5, 6, 10$.
Key Words: quaternary quadratic forms; theta functions; "The 290-theorem".
MSC(2010) 14K25; 11E25

1 Introduction

Throughout this paper, set $\mathbb{N} = \{1, 2, 3, \ldots\}$ and $\mathbb{N}_0 = \{0, 1, 2, 3, \ldots\}$. $\mathbb{Z}$ denotes the set of rational integers. In addition, the triangular numbers are $t_x = x(x+1)/2,\ (x \in \mathbb{N}_0)$ and the squares are $y^2,\ (y \in \mathbb{Z})$. Furthermore, a, b and c are fixed positive integers with $1 \leq a \leq b$.

Lagrange proved that every $n \in \mathbb{N}_0$ is a sum of four squares. Noted is that Jacobi [6] showed this fact using the elliptic function theory.

Ramanujan [7] determined all the positive integers a, b, c and d such that every $n \in \mathbb{N}_0$ is represented as $ax^2 + by^2 + cz^2 + du^2$ with $x, y, z, u \in \mathbb{Z}$. He proved that there exist fifty-four such quadruples (a, b, c, d) with $1 \leq a \leq b \leq c \leq d$. A proof of this fact is contained in Dickson [4].

The aim of this paper is to determine all the positive integers a, b and c with $a \leq b$ such that every $n \in \mathbb{N}_0$ can be represented as

$$f_c^{a,b}(x,y,z,w) = ax^2 + by^2 + c(z^2 + zw + w^2)$$

with $x, y, z, w \in \mathbb{Z}$. Furthermore, we prove that $f_c^{a,b}$ can represent all the nonnegative integers if it represents $n = 1, 2, 3, 5, 6, 10$. Our main theorem is as follows:

Theorem 1.1. *For positive integers a, b and c with $a \leq b$, set*

$$f_c^{a,b}(x,y,z,w) = ax^2 + by^2 + c(z^2 + zw + w^2),\ (x, y, z, w \in \mathbb{Z}).$$

(1) *$f_c^{a,b}$ can represent any nonnegative integers if and only if*

$$(a,b,c) = \begin{cases} (1,b,1) & (b=1,2,3,4,5,6), \\ (2,b,1) & (b=2,3,4,5,6,7,8,9,10), \\ (1,b,2) & (b=1,2,3,4,5), \\ (1,2,3), \\ (1,2,4). \end{cases}$$

(2) *If $f_c^{a,b}$ represents $n = 1,2,3,5,6,10$, then it can represent all the nonnegative integers.*

Remarks

Theorem 1.1 (2) is the restriction of "The 290-theorem" to the forms $ax^2 + by^2 + c(z^2 + zw + w^2)$. Bhargava and Hanke [1] proved "The 290-theorem".

Theorem 1.2. ("The 290-theorem") *If a positive-definite quadratic form with integer coefficients represents the 29 integers*

$$1,2,3,5,6,7,10,13,14,15,17,19,21,22,23,26,$$
$$29,30,31,34,35,37,42,58,93,110,145,203,290,$$

then it represents all positive integers.

For the proof of Theorem 1.1, following Ramanujan, we introduce

$$\varphi(q) = \sum_{n\in\mathbb{Z}} q^{n^2},\ \psi(q) = \sum_{n=0}^{\infty} q^{\frac{n(n+1)}{2}},\ a(q) = \sum_{m,n\in\mathbb{Z}} q^{m^2+mn+n^2},\ (q\in\mathbb{C},\ |q|<1),$$

and use the following identities:

$$a(q) = a(q^4) + 6q\psi(q^2)\psi(q^6). \tag{1.1}$$
$$\varphi(q)\varphi(q^3) = a(q^4) + 2q\psi(q^2)\psi(q^6). \tag{1.2}$$

For the proof of these formulas, see Berndt [2, pp. 232] and Hirschhorn et al. [5].

Acknowledgments

We are grateful to Professor Williams for informing us the "The 290-theorem" and, for his useful suggestions.

2 Notations and preliminaries

For the fixed positive integers a, c and each $n \in \mathbb{N}_0$, we define

$$f_c^a(x,y,z) = ax^2 + c(y^2 + yz + z^2),$$

and

$$A_c^a(n) = \sharp\left\{(x,y,z) \in \mathbb{Z}^3 \,|\, n = f_c^a(x,y,z)\right\}.$$

Moreover, for the fixed positive integers α, β, γ and each $n \in \mathbb{N}_0$, we set

$$\begin{aligned} r_{\alpha,\beta}(n) =& \sharp\left\{(x,y) \in \mathbb{Z}^2 \,|\, n = \alpha x^2 + \beta y^2\right\}, \\ t_{\alpha,\beta}(n) =& \sharp\left\{(x,y) \in \mathbb{N}_0^2 \,|\, n = \alpha t_x + \beta t_y\right\}, \\ r_{\alpha,\beta,\gamma}(n) =& \sharp\left\{(x,y,z) \in \mathbb{Z}^3 \,|\, n = \alpha x^2 + \beta y^2 + \gamma z^2\right\}, \\ m_{\alpha\text{-}\beta,\gamma}(n) =& \sharp\left\{(x,y,z) \in \mathbb{Z} \times \mathbb{N}_0^2 \,|\, n = \alpha x^2 + \beta t_y + \gamma t_z\right\}. \end{aligned}$$

3 The case where $c = 1$

Lemma 3.1. Suppose that $a \leq b$, $c = 1$ and $f_c^{a,b}(x,y,z,w)$, $(x,y,z,w \in \mathbb{Z})$ represent any nonnegative integers $n \in \mathbb{N}_0$. Then, $a = 1, 2$.

Proof. The lemma follows from the fact that $n = 2$ cannot be written as $z^2 + zw + w^2$ with $z, w \in \mathbb{Z}$. □

3.1 The case where $a = 1$

We use the following result of Dickson [4, pp. 112-113]:

Lemma 3.2. *$n \in \mathbb{N}_0$ can be written as $x^2 + y^2 + 3z^2$ with $x, y, z \in \mathbb{Z}$ if and only if $n \neq 9^k(9l + 6)$, $(k, l \in \mathbb{N}_0)$.*

By Lemma 3.2, we have the following proposition:

Proposition 3.3. *$n \in \mathbb{N}_0$ can be written as $x^2 + (y^2 + yz + z^2)$ with $x, y, z \in \mathbb{Z}$ if and only if $n \neq 9^k(9l + 6)$, $(k, l \in \mathbb{N}_0)$.*

Proof. Multiplying both sides of equations (1.1) and (1.2) by $\varphi(q)$, we have

$$\varphi(q)a(q) = \varphi(q)a(q^4) + 6q\varphi(q)\psi(q^2)\psi(q^6), \tag{3.1}$$

$$\varphi(q)^2\varphi(q^3) = \varphi(q)a(q^4) + 2q\varphi(q)\psi(q^2)\psi(q^6). \tag{3.2}$$

which implies that

$$\sum_{n=0}^{\infty} A_1^1(n)q^n = \sum_{n=0}^{\infty} A_4^1(n)q^n + 6q\sum_{N=0}^{\infty} m_{1\text{-}2,6}(N)q^N,$$
$$\sum_{n=0}^{\infty} r_{1,1,3}(n)q^n = \sum_{n=0}^{\infty} A_4^1(n)q^n + 2q\sum_{N=0}^{\infty} m_{1\text{-}2,6}(N)q^N.$$

Therefore, it follows that

$$\begin{aligned} n \neq 9^k(9l+6),\ (k,l \in \mathbb{N}_0) &\Longleftrightarrow r_{1,1,3}(n) > 0 \\ &\Longleftrightarrow A_4^1(n) > 0 \text{ or } m_{1\text{-}2,6}(n-1) > 0, \\ &\Longleftrightarrow A_1^1(n) > 0, \end{aligned}$$

which proves the proposition. □

By Proposition 3.3, we obtain the following theorem:

Theorem 3.4. *$f_1^{1,b}$ can represent any nonnegative integers $n \in \mathbb{N}_0$ if and only if $b = 1, 2, 3, 4, 5, 6$.*

Proof. The "only if" direction follows from Proposition 3.3. In order to establish the "if" direction, we have only to prove that $f_1^{1,b}$ represents $n = 9l + 6,\ (l \in \mathbb{N}_0)$.

When $b = 1, 2, 3, 4, 5$, taking $y = 1$, we have

$$n - by^2 = 9l + 6 - b \cdot 1^2 \equiv 5, 4, 3, 2, 1 \bmod 9,$$

which can be written as $x^2 + (z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$.

Suppose that $b = 6$. If $n = 6, 15$, or $l \equiv 2, 8 \bmod 9$, taking $y = 1$, we obtain

$$n - 6 \cdot 1^2 = 9l,\ (l = 0, 1 \text{ or } l \equiv 2, 8 \bmod 9),$$

which can be represented as $x^2 + (z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$.

If $l \geq 2$ and $l \not\equiv 2, 8 \bmod 9$, taking $y = 2$, we have

$$n - 6 \cdot 2^2 = 9l + 6 - 24 = 9(l - 2),$$

which can be written as $x^2 + (z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$. □

3.2 The case where $a = 2$

We use the following result of Dickson [3]:

Lemma 3.5.

(1) *$n \in \mathbb{N}_0$ can be written as $x^2 + 2y^2 + 3z^2$ with $x, y, z \in \mathbb{Z}$ if and only if $n \neq 4^k(16l + 10)$, $(k, l \in \mathbb{N}_0)$.*

(2) *$n \in \mathbb{N}_0$ can be written as $x^2 + 2(y^2 + yz + z^2)$ with $x, y, z \in \mathbb{Z}$ if and only if $n \neq 4^k(8l + 5)$, $(k, l \in \mathbb{N}_0)$.*

By Lemma 3.5, we have the following proposition:

Proposition 3.6. *$n \in \mathbb{N}_0$ can be written as $2x^2 + (y^2 + yz + z^2)$ with $x, y, z \in \mathbb{Z}$ if and only if $n \neq 4^k(16l + 10)$, $(k, l \in \mathbb{N}_0)$.*

Proof. Multiplying both sides of equations (1.1) and (1.2) by $\varphi(q^2)$, we have

$$\varphi(q^2)a(q) = \varphi(q^2)a(q^4) + 6q\varphi(q^2)\psi(q^2)\psi(q^6), \tag{3.3}$$
$$\varphi(q)\varphi(q^2)\varphi(q^3) = \varphi(q^2)a(q^4) + 2q\varphi(q^2)\psi(q^2)\psi(q^6). \tag{3.4}$$

which implies that

$$\sum_{n=0}^{\infty} A_1^2(n)q^n = \sum_{N=0}^{\infty} A_2^1(N)q^{2N} + 6q\sum_{N=0}^{\infty} m_{1\text{-}1,3}(N)q^{2N},$$
$$\sum_{n=0}^{\infty} r_{1,2,3}(n)q^n = \sum_{N=0}^{\infty} A_2^1(N)q^{2N} + 2q\sum_{N=0}^{\infty} m_{1\text{-}1,3}(N)q^{2N}.$$

Suppose that n is even and $n = 2N$. Therefore, it follows that

$$\begin{aligned} n \neq 4^k(16l + 10),\ (k, l \in \mathbb{N}_0) &\iff r_{1,2,3}(n) > 0 \\ &\iff A_2^1(N) > 0 \\ &\iff A_1^2(n) > 0. \end{aligned}$$

Suppose that n is odd and $n = 2N + 1$. Therefore, it follows that

$$\begin{aligned} n \neq 4^k(16l + 10),\ (k, l \in \mathbb{N}_0) &\iff r_{1,2,3}(n) > 0 \\ &\iff m_{1\text{-}1,3}(N) > 0 \\ &\iff A_1^2(n) > 0. \end{aligned}$$

□

By Proposition 3.6, we obtain the following theorem:

Theorem 3.7. *Suppose that $2 \leq b$. Then, $f_1^{2,b}$ can represent any nonnegative integers $n \in \mathbb{N}_0$ if and only if $b = 2, 3, 4, 5, 6, 7, 8, 9, 10$.*

Proof. The "only if" direction follows from Proposition 3.6. In order to establish the "if" direction, we have only to prove that $f_1^{2,b}$ represents $n = 16l + 10$, $(l \in \mathbb{N}_0)$.

When $b \neq 2, 10$, taking $y = 1$, we have

$$n - by^2 = 16l + 10 - b \cdot 1^2 \not\equiv 0, 8, 10 \bmod 16,$$

which can be written as $2x^2 + (z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$.

When $b = 2$, taking $y = 2$, we obtain

$$n - 2 \cdot 2^2 = 16l + 10 - 8 = 16l + 2,$$

which can be represented as $2x^2 + (z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$.

We last suppose that $b = 10$. If $n = 10, 26$, taking $y = 1$, we obtain

$$n - 10 \cdot 1^2 = 0, 16,$$

which can be written as $2x^2 + (z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$.

If $l \geq 2$, taking $y = 2$, we have

$$n - 10 \cdot 2^2 = 16l + 10 - 40 = 16(l - 2) + 2,$$

which can be represented as $2x^2 + (z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$. □

3.3 Summary

By Theorems 3.4, 3.7 and their proofs, we obtain

Theorem 3.8. *Let a, b be positive integers with $a \leq b$.*

(1) *$f_1^{a,b}$ can represent any nonnegative integers if and only if (a, b) are given by*

$$(a, b) = \begin{cases} (1, b) & (b = 1, 2, 3, 4, 5, 6), \\ (2, b) & (b = 2, 3, 4, 5, 6, 7, 8, 9, 10). \end{cases}$$

(2) *If $f_1^{a,b}$ represents $n = 1, 2, 6, 10$, it can represent all the nonnegative integers.*

Furthermore, we obtain the following theorem:

Theorem 3.9. *For fixed positive integers, a, c, set*

$$f_c^a(x, y, z) = ax^2 + c(y^2 + yz + z^2) \text{ with } x, y, z \in \mathbb{Z}.$$

There exists no positive integers, a, c such that f_c^a can represent every natural number.

Proof. Suppose that there exist such positive integers, a, c. Taking $n = 1$, we have $a = 1$ or $c = 1$.

Suppose that $a = 1$. The choice $n = 2$ implies that $c = 1, 2$. On the other hand, if $(a, c) = (1, 1), (1, 2)$, by Proposition 3.3 and Lemma 3.5, we see that there exist positive integers which cannot be expressed by f_1^1 or f_2^1, which is contradiction.

Suppose that $c = 1$. The choice $n = 2$ implies that $a = 1, 2$. If $(a, c) = (1, 1), (2, 1)$, by Propositions 3.3 and 3.6, we see that there exist positive integers which cannot be expressed by f_1^1 or f_2^1, which is contradiction. □

Remark

In [4, pp.104], Dickson proved that there exist no positive integers, a, b, c, such that $ax^2 + by^2 + cz^2$, $(x, y, z \in \mathbb{Z})$ can represent all positive integers.

4 The case where $c = 2$

Noting that $a = 1$ if $f_2^{a,b}$ can represent any nonnegative integers, by Lemma 3.5 (2), we obtain the following theorem:

Theorem 4.1. *Suppose that $1 \leq a \leq b$. Then, $f_2^{a,b}$ can represent any nonnegative integers $n \in \mathbb{N}_0$ if and only if $a = 1$ and $b = 1, 2, 3, 4, 5$.*

Proof. The "only if" direction follows from Lemma 3.5 (2). In order to establish the "if" direction, we have only to prove that $f_2^{1,b}$ represents $n = 8l + 5$, $(l \in \mathbb{N}_0)$.

We first treat the case where $b = 1$. Taking $y = 2$, we obtain

$$n - 1 \cdot 2^2 = 8l + 5 - 4 = 8l + 1,$$

which can be represented as $x^2 + 2(z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$.

When $b = 2, 3, 4$, taking $y = 1$, we have

$$n - by^2 = 8l + 5 - b \cdot 1^2 \equiv 3, 2, 1 \bmod 8,$$

which can be written as $x^2 + 2(z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$.

We last suppose that $b = 5$. If $l = 0, 1$, taking $y = 1$, we obtain

$$n - 5 \cdot 1^2 = 8l + 5 - 5 = 0, 8,$$

which can be represented as $x^2 + 2(z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$.

If $l \geq 2$, taking $y = 2$, we have

$$n - 5 \cdot 2^2 = 8l + 5 - 20 = 8(l - 2) + 1,$$

which can be written as $x^2 + 2(z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$. □

By the proof of Theorem 4.1, we obtain

Theorem 4.2. *If $f_2^{a,b}$ represents $n = 1, 5$, it can represent all the nonnegative integers.*

5 The case where $c = 3$

We use the result of Dickson [4, pp. 112-113]:

Lemma 5.1. *$n \in \mathbb{N}_0$ can be written as $x^2 + 3y^2 + 9z^2$ with $x, y, z \in \mathbb{Z}$ if and only if $n \neq 3l + 2,\ 9^k(9l + 6),\ (k, l \in \mathbb{N}_0)$.*

By Lemma 5.1, we have the following proposition:

Proposition 5.2. *$n \in \mathbb{N}_0$ can be written as $x^2 + 3(y^2 + yz + z^2)$ with $x, y, z \in \mathbb{Z}$ if and only if $n \neq 3l + 2,\ 9^k(9l + 6),\ (k, l \in \mathbb{N}_0)$.*

Proof. Replacing q by q^3 in equations (1.1) and (1.2), we have

$$\begin{aligned} a(q^3) =& a(q^{12}) + 6q^3\psi(q^6)\psi(q^{18}), \\ \varphi(q^3)\varphi(q^9) =& a(q^{12}) + 2q^3\psi(q^6)\psi(q^{18}). \end{aligned}$$

Multiplying both sides of these equations by $\varphi(q)$, we have

$$\varphi(q)a(q^3) = \varphi(q)a(q^{12}) + 6q^3\varphi(q)\psi(q^6)\psi(q^{18}), \tag{5.1}$$

$$\varphi(q)\varphi(q^3)\varphi(q^9) = \varphi(q)a(q^{12}) + 2q^3\varphi(q)\psi(q^6)\psi(q^{18}). \tag{5.2}$$

which implies that

$$\begin{aligned} \sum_{n=0}^{\infty} A_3^1(n)q^n =& \sum_{n=0}^{\infty} A_{12}^1(n)q^n + 6q^3 \sum_{N=0}^{\infty} m_{1\text{-}6,18}(N)q^N, \\ \sum_{n=0}^{\infty} r_{1,3,9}(n)q^n =& \sum_{n=0}^{\infty} A_{12}^1(n)q^n + 2q^3 \sum_{N=0}^{\infty} m_{1\text{-}6,18}(N)q^N. \end{aligned}$$

Therefore, it follows that

$$\begin{aligned} n \neq 3l+2,\ 9^k(9l+6),\ (k,l \in \mathbb{N}_0) &\Longleftrightarrow r_{1,3,9}(n) > 0 \\ &\Longleftrightarrow A^1_{12}(n) > 0 \,\text{or}\, m_{1\text{-}6,18}(n-3) > 0, \\ &\Longleftrightarrow A^1_3(n) > 0, \end{aligned}$$

which proves the proposition. □

Noting that $a = 1$ if $f_3^{a,b}$ can represent any nonnegative integers, we obtain the following theorem:

Theorem 5.3. *$f_3^{a,b}$ can represent any nonnegative integers $n \in \mathbb{N}_0$ if and only if $a = 1$ and $b = 2$.*

Proof. Let us first prove the "only if" direction. Proposition 5.2 implies that $b = 1, 2$. Assume that $b = 1$ and

$$n = 6 = x^2 + y^2 + 3(z^2 + zw + w^2) \text{ with } x, y, z, w \in \mathbb{Z},$$

which implies that $z^2 + zw + w^2 = 1, \neq 0, 2$. It then follows that $x^2 + y^2 = 3$, which is impossible.

In order to establish the "if" direction, we have only to prove that $f_3^{1,b}$ can represent $n = 3l+2,\ 9l+6,\ (l \in \mathbb{N}_0)$.

Suppose that $n = 3l + 2$. We first treat the case where $l \equiv 0 \bmod 3$. It is obvious that $f_3^{1,2}$ can represent $n = 2$. When $l \geq 1$ and $l \equiv 0 \bmod 3$, we obtain

$$n - 2 \cdot 2^2 = 3l + 2 - 8 = 3(l-2),$$

which can be written as $x^2 + 3(z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$.

When $l \equiv 1 \bmod 3$, taking $y = 1$, we have

$$n - 2y^2 = 3l + 2 - 2 = 3l,$$

which can be written as $x^2 + 3(z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$.

Assume that $l \equiv 2 \bmod 3$. For $l = 2, 5, 8$, taking $y = 2$, we have

$$n - 2 \cdot 2^2 = 3l + 2 - 2 \cdot 2^2 = 0, 9, 9 \cdot 2,$$

which can be represented as $x^2 + 3(z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$. If $l \geq 11$ and $l = 3L + 2,\ (L \in \mathbb{N}_0)$, taking $y = 4$, we have

$$n - 2 \cdot 4^2 = 3l + 2 - 2 \cdot 4^2 = 3\{3(L-3) + 1\},$$

which can be written as $x^2+3(z^2+zw+w^2)$ with $x,z\in\mathbb{Z}$.

We last suppose that $n=9l+6$. Taking $y=1$, we have

$$n-2y^2=9l+6-2=9l+4\equiv 1 \bmod 3,$$

which can be represented as $x^2+3(z^2+zw+w^2)$ with $x,z\in\mathbb{Z}$. □

By the proof of Theorem 5.3, we obtain

Theorem 5.4. *If $f_3^{a,b}$ represents $n=1,2,6$, it can represent all the nonnegative integers.*

6 The case where $c=4$

We use the following result of Dickson [4, pp. 112-113]:

Lemma 6.1. *$n\in\mathbb{N}_0$ can be written as $x^2+4y^2+12z^2$ with $x,y,z\in\mathbb{Z}$ if and only if $n\neq 4l+2,\ 4l+3,\ 9^k(9l+6),\ (k,l\in\mathbb{N}_0)$.*

By Lemma 6.1, we have the following proposition:

Proposition 6.2. *$n\in\mathbb{N}_0$ can be written as $x^2+4(y^2+yz+z^2)$ with $x,y,z\in\mathbb{Z}$ if and only if $n\neq 4l+2,\ 4l+3,\ 9^k(9l+6),\ (k,l\in\mathbb{N}_0)$.*

Proof. Replacing q by q^4 in equations (1.1) and (1.2), we have

$$\begin{aligned} a(q^4) =& a(q^{16})+6q^4\psi(q^8)\psi(q^{24}),\\ \varphi(q^4)\varphi(q^{12}) =& a(q^{16})+2q^4\psi(q^8)\psi(q^{24}). \end{aligned}$$

Multiplying both sides of these equations by $\varphi(q)$, we have

$$\varphi(q)a(q^4) =\varphi(q)a(q^{16})+6q^4\varphi(q)\psi(q^8)\psi(q^{24}), \tag{6.1}$$

$$\varphi(q)\varphi(q^4)\varphi(q^{12}) =\varphi(q)a(q^{16})+2q^4\varphi(q)\psi(q^8)\psi(q^{24}). \tag{6.2}$$

which implies that

$$\begin{aligned} \sum_{n=0}^{\infty}A_4^1(n)q^n =& \sum_{n=0}^{\infty}A_{16}^1(n)q^n+6q^4\sum_{N=0}^{\infty}m_{1\text{-}8,24}(N)q^N,\\ \sum_{n=0}^{\infty}r_{1,4,12}(n)q^n =& \sum_{n=0}^{\infty}A_{16}^1(n)q^n+2q^4\sum_{N=0}^{\infty}m_{1\text{-}8,24}(N)q^N. \end{aligned}$$

Therefore, it follows that

$$\begin{aligned} n \neq 4l+2,\ 4l+3,\ 9^k(9l+6),\ (k,l \in \mathbb{N}_0) &\Longleftrightarrow r_{1,4,12}(n) > 0 \\ &\Longleftrightarrow A_{16}^1(n) > 0 \,\text{or}\, m_{1\text{-}8,24}(n-4) > 0, \\ &\Longleftrightarrow A_4^1(n) > 0, \end{aligned}$$

which proves the proposition. □

Noting that $a = 1, b = 2$ if $f_4^{a,b}$ can represent any nonnegative integers, we obtain the following theorem:

Theorem 6.3. *$f_4^{a,b}$ can represent any nonnegative integers $n \in \mathbb{N}_0$ if and only if $a = 1$ and $b = 2$.*

Proof. The "only if" direction is obvious. In order to establish the "if" direction, we have only to prove that $f_4^{1,2}$ represents $n = 4l+2,\ 4l+3,\ 9l+6,\ (l \in \mathbb{N}_0)$.

Suppose that $n = 4l+2$. If $l \not\equiv 0, 6 \bmod 9$, taking $y = 1$, we have

$$n - 2y^2 = 4l + 2 - 2 = 4l,$$

which can be represented as $x^2 + 4(z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$. If $l \equiv 0, 6 \bmod 9$, taking $y = 3$, we obtain

$$n - 2y^2 = 4l + 2 - 2 \cdot 3^2 = 4(l-4),$$

which can be represented as $x^2 + 4(z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$.

Suppose that $n = 4l+3$. If $l \not\equiv 2, 8 \bmod 9$, taking $y = 1$, we have

$$n - 2y^2 = 4l + 3 - 2 = 4l + 1 \not\equiv 0, 6 \bmod 9,$$

which can be represented as $x^2 + 4(z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$. If $l \equiv 2, 8 \bmod 9$, taking $y = 3$, we obtain

$$n - 2y^2 = 4l + 3 - 2 \cdot 3^2 = 4(l-4) + 1 \equiv 2, 8 \bmod 9,$$

which can be represented as $x^2 + 4(z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$. It is easy to check that $f_4^{1,2}$ can represent $n = 11$.

Suppose that $n = 9l+6$. If $l \equiv 0 \bmod 4$ and $l = 4L,\ (L \in \mathbb{N}_0)$, taking $y = 5$, we have

$$n - 2y^2 = 9l + 6 - 2 \cdot 5^2 = 4\{9(L-2) + 7\},$$

which can be written as $x^2 + 4(z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$. It is easy to check that $f_4^{1,2}$ can represent $n = 6, 42$.

If $l \equiv 1 \bmod 4$, taking $y = 1$, we have

$$n - 2y^2 = 9l + 6 - 2 \cdot 1^2 = 9l + 4 \equiv 1 \bmod 4,$$

which can be represented as $x^2 + 4(z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$.

If $l \equiv 2 \bmod 4$ and $l = 4L + 2$, $(L \in \mathbb{N}_0)$, taking $y = 4$, we have

$$n - 2y^2 = 9l + 6 - 2 \cdot 4^2 = 4(9L - 2),$$

which can be written as $x^2 + 4(z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$. It is easy to check that $f_4^{1,2}$ can represent $n = 24$.

If $l \equiv 3 \bmod 4$, taking $y = 2$, we have

$$n - 2y^2 = 9l + 6 - 2 \cdot 2^2 = 9l - 2 \equiv 1 \bmod 4,$$

which can be represented as $x^2 + 4(z^2 + zw + w^2)$ with $x, z \in \mathbb{Z}$. □

By the proof of Theorem 6.3, we obtain

Theorem 6.4. *If $f_4^{a,b}$ represents $n = 1, 2, 3$, it can represent all the nonnegative integers.*

7 The case where $c \geq 5$

7.1 The case where $c = 5$

Theorem 7.1. *There exist no positive integers a, b such that $1 \leq a \leq b$ and $f_5^{a,b}$ can represent any nonnegative integers $n \in \mathbb{N}_0$, where*

$$f_5^{a,b}(x, y, z, w) = ax^2 + by^2 + 5(z^2 + zw + w^2).$$

Proof. Suppose that there exist such positive integers a and b. Considering $n = 1$, we have $a = 1$. Taking $n = 2$, we obtain $b = 1, 2$. Considering $n = 3$, we have $(a, b) = (1, 2)$.

Take $n = 10$. Then, n cannot be written as $x^2 + 2y^2$ with $x, y \in \mathbb{Z}$, which implies that $z^2 + zw + w^2 = 1$, because 2 cannot be represented as $z^2 + zw + w^2$ with $z, w \in \mathbb{Z}$. Therefore, it follows that $5 = 10 - 5 \cdot 1$ can be written as $x^2 + 2y^2$ with $x, y \in \mathbb{Z}$, which is impossible. □

7.2 The case where $c \geq 6$

Theorem 7.2. *There exist no positive integers a, b, c such that $1 \leq a \leq b$, $c \geq 6$ and $f_c^{a,b}$ can represent any nonnegative integers $n \in \mathbb{N}_0$.*

Proof. Suppose that there exist such positive integers a, b and c. Considering $n = 1$, we have $a = 1$. Taking $n = 2$, we obtain $b = 1, 2$. Considering $n = 3$, we have $(a, b) = (1, 2)$. On the other hand, $n = 5$ cannot be written as $x^2 + 2y^2$ with $x, y \in \mathbb{Z}$. □

8 Proof of Theorem 1.1

Proof. Theorem 1.1 (1) follows from Theorems 3.8, 4.1, 5.3, 6.3, 7.1 and 7.2. Theorem 1.1 (2) follows from Theorems 3.8, 4.2, 5.4, 6.4, 7.1 and 7.2. □

References

[1] M. Bhargava and J. Hanke, *Universal Quadratic Forms and the 290-Theorem,* (preprint)

[2] B. C. Berndt, *Ramanujan's notebooks. Part III.* Springer-Verlag, New York, (1991).

[3] L. E. Dickson, *Integers represented by positive ternary quadratic forms,* Bull. Amer. Math. Soc. **33** 63-70 (1927).

[4] L. E. Dickson, *Modern Elementary Theory of Numbers,* University of Chicago Press, Chicago, (1939).

[5] M. Hirschhorn, F. Garvan, and J. Borwein, *Cubic analogues of the Jacobian theta function* $\theta(z, q)$, Canad. J. Math. **45** 673-694 (1993).

[6] C. G. J. Jacobi, *Fundamenta Nova Theoriae Functionum Ellipticarum,* Bornträger, Regiomonti, (1829).

[7] S. Ramanujan, *On the expression of a number in the form* $ax^2 + by^2 + cz^2 + du^2$, Proc. Cambridge Philos. Soc. **19** 11-21 (1917).